Children's Healthcare of Atlanta

My asthma action plan

octor's name:		Signature:	
octor's phone #:	•		
Controller medicines	How much to take	How often	Other instructions
		times per day	☐ Gargle or rinse mouth after use
		times per day	
,		times per day	
Quick-relief medicines	How much to take	How often	Other instructions
	☐ 2 puffs ☐ 4-6 puffs ☐ 1 nebulizer treatment	Take ONLY as needed (see below.— starting in Yellow Zone or before excercise)	NOTE: If you need this medicine more than 2 days a week, call your doctor.
Asthma triggers (check all that □ Exercise □ Change in to □ Pollens □ Respiratory	emperature	☐ Strong emotions	odors or fumes
Special ins	tructions when I am		ma symptoms every day:
 No coughing, wheezing, tightness, shortness of briddering the day or night Can go to school and play Be careful. Coughing, wheezing, che tightness, shortness of bridghtness, shortness of bridghtness of bridghtness, shortness of bridghtness, shortness of bridghtness of bridghtness, shortness of bridghtness of bridghtnes	eath y	Avoid trigg (See above) CAUTION. Cont Take quick relief within one h Continue us needed. Ca	rcise, take puff(s) of ers that make my asthma worse inue taking my controller medicines every day. puffs or nebulizer treatment(s) of medicine. If I am not back in the Green Zone nour, then I should: sing quick relief medicine every 4 hours as Il provider if not improving in days.
Ask for help. • Very short of breath • Continual coughing • Skin between ribs is pullir • Difficulty speaking withou out of breath • Quick-relief medicines had not helped • Symptoms same or worse	at running ve	Take quick-minutes and	ERT! Get help! relief medicine: puffs every d get help immediately.

314 Children's Healthcare of Atlanta Inc. All rights reserved. MED 95